

P1

Is ankle-brachial pressure index sufficient test for indication of percutaneous transluminal angioplasty in patients with diabetic foot ulcers?

V. Woskova, A. Jirkovska, R. Bem, M. Dubsky, K. Cechova, J. Peregrin, B. Koznar
Diabetes Center IKEM, Prague, Czech Republic

Background: The decision to perform vascular intervention is often based on Doppler ankle/brachial index (ABI). The ABI <0.9 indicates peripheral arterial disease (PAD) according to International Consensus on the Diabetic Foot but medial arterial sclerosis may be the confounding factor of this cut-point. The relationship between ABI prior to percutaneous transluminal angioplasty (PTA) and PTA outcomes is not proved enough. The aim of our study was to compare short- and long-term effect of PTA between diabetic patients with ABI ≥ 0.9 and with ABI <0.9 . **Patients and methods:** The retrospective study was conducted to evaluate 49 lower limbs in 47 diabetic patients (Type1/Type2 DM 9/38, mean age 63 ± 15 years, mean diabetes duration 20 ± 11 years) treated at a diabetic foot clinic for diabetic foot syndrome (DFS) during the period of 2005-2007, in which non-invasive vascular assessment was documented. The mean follow up period was 10 (6-12) month. Indication for PTA were non-healing diabetic foot ulcers. In the first group lower limbs with ABI ≥ 0.9 were included, in the second group lower limbs with ABI < 0.9 were included. The evaluation of short-term effect of PTA was based on transcutaneous oxygen tension (TcPO₂) measurement before and 3-6 days after PTA; evaluation of long-term effect was based on clinical observation: positive effect was defined as ulcer closure or below-the ankle amputation healed and healing failure was defined as non-healed ulcer or above-the-ankle amputation. **Results:** In the first group (ABI ≥ 0.9) 23 lower limbs with mean ankle pressure 172 ± 35 mmHg and mean ABI $1,26 \pm 0,24$; in the second group (ABI <0.9) 26 lower limbs with mean ankle pressure 83 ± 29 mmHg and mean ABI $0,58 \pm 0,19$ were evaluated. Using Wagner classification both groups were comparable as to severity of lesions. Both groups didn't differ significantly in the mean TcPO₂ before the PTA (23.1 ± 15.9 vs 23.6 ± 16.3 mmHg, NS). There was significant increase of TcPO₂ after PTA both in the first ($\Delta 12.9 \pm 18.4$ mmHg, $p=0.0035$) and the second group ($\Delta 13.9 \pm 13.9$ mmHg, $p<0.001$) with no significant differences between both groups. No significant difference between first and second group in clinical outcomes during the follow up period was observed: ulcer healing in 18/23 vs 17/26 (78% vs 65%, NS) and healing failure in 5/23 vs 9/26 (22% vs 35%, NS). **Conclusions:** Our study demonstrated no significant differences in short-term and long-term effect of PTA between patients with ABI ≥ 0.9 and in patients with ABI < 0.9 . We concluded, that ABI is not sufficient criterion for PTA indication in patients with non-healing diabetic foot ulcers and ABI ≥ 0.9 cannot exclude positive effect of PTA on ulcer healing. This study was supported by grant MZO 00023001 from the Ministry of Health of the Czech Republic.